

# Onslaught®

## MICROENCAPSULATED INSECTICIDE

- Kills for up to 12 months (1 year)
- Kills German cockroaches for up to 12 months (1 year) on indoor, unsealed concrete and/or cement surfaces only

### ACTIVE INGREDIENT:

(S)-cyano (3-phenoxyphenyl) methyl-(S)-4-chloro-alpha-(1-methylethyl) benzenoacetate .....	6.40%
OTHER INGREDIENTS* .....	93.60%
	100.00%

\*Contains petroleum distillates

KEEP OUT OF REACH OF CHILDREN

## CAUTION

See reverse for first aid and precautionary statements.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Remove pets and birds and cover fish aquariums before spraying.

For control of insects, indoors and outdoors, in food and non-food areas such as, but not limited to: homes, schools, warehouses, office buildings, apartment buildings, theatres, hotels, industrial buildings, motels, kennels, livestock housing, food processing plants, food service establishments, restaurants, supermarkets and grocery stores, transportation equipment, truck trailers, railroad cars, and food manufacturing and warehousing establishments. Also for use on backyards, lawns, trees, ornamental landscaping, recreational areas, parks and athletic fields.

### INSECTS CONTROLLED:

Ants  
Aphids  
Asian Lady Beetles  
Bed Bugs  
Beetles  
Boxelder Bugs  
Carpet Beetles  
Centipedes  
Cockroaches  
Crickets  
Dog Ticks  
Earwigs  
Fire Ants  
Firebrats  
Fleas  
Mealy Bugs  
Midges  
Millipedes  
Mites  
Pillbugs  
Psocids  
Scales  
Silverfish  
Spiders  
Springtails  
Swarming Termites  
Ticks  
Waterbugs

Fruit Flies  
Fungus Gnats  
Gnats  
Horn Flies  
Houseflies  
Mosquitoes  
Small Flying Moths  
Whiteflies

### BITING AND STINGING PESTS:

Bed Bugs  
Bees  
Biting Flies  
Chiggers  
Deer Flies  
Deer Ticks  
Dog Ticks  
Fire Ants  
Fleas  
Gnats  
Hornets  
Lice  
Mosquitoes  
Scorpions  
Spiders  
Ticks  
Wasps  
Yellow Jackets

### STORED PRODUCT PESTS INCLUDING:

Angoumois Grain Moths  
Ants  
Cadellés  
Cigarette Beetles  
Cockroaches  
Confused Flour Beetles

Dark Mealworms  
Dried Fruit Beetles  
Drug Store Beetles  
Flat Grain Beetles  
Fruit Flies  
Grain Mites  
Grain Moths  
Granary Weevils  
Indian Meal Moths  
Lesser Grain Borers  
Maize Weevils  
Meal Moth Larvae  
Mediterranean Flour Moths  
Merchant Grain Beetles  
Red Flour Beetles  
Rice Weevils  
Rusty Grain Beetles  
Saw-toothed Grain Beetles  
Skippers  
Spider Beetles  
Spider Mites  
Square-necked Grain Beetles  
Tobacco Moths  
Yellow Mealworms

### LIVESTOCK PREMISE PESTS INCLUDING:

Bed Bugs  
Carrion Beetles  
Chiggers  
Darkling Beetles (lesser meal worm)  
Deer Flies  
Face Flies  
Fleas  
Flies  
Hide Beetles

Horn Flies  
Horse Flies  
Lice  
Litter Beetles  
Mites  
Mosquitoes  
Stable Flies

### WOOD DESTROYING PESTS INCLUDING:

Carpenter Ants  
Carpenter Bees  
Deathwatch Beetles  
Furniture Beetles  
Old House Borers  
Powder Post Beetles  
Round-headed House Borers  
Swarming Termites

### PLANT PESTS INCLUDING:

Alfalfa Caterpillars  
American Plum Borers  
Annual Bluegrass Weevils  
Aphids  
Apple Maggots  
Armyworms  
Artichoke Plume Moths  
Bagworms  
Balsam Woolly Adelgids  
Beet Armyworms  
Billbugs  
Blueberry Spanworms  
Cherry Fruit Flies  
Cherry Fruitworms  
Chinch Bugs  
Codling Moths  
Cowpea Curculios

Cranberry Fruitworms  
Cranberry Weevils  
Cucumber Beetles  
Cutworms  
Diamondback Moths  
Elm Leaf Beetles  
European Pine Sawflies  
Fall Webworms  
Filbert Worms  
Flea Beetles  
Grasshoppers  
Green Cloverworms  
Green Fruit Worms  
Gypsy Moth Caterpillars  
Hickory Shuckworms  
Imported Cabbageworms  
Japanese Beetles  
Lace Bugs  
Leaf Feeding Caterpillars  
Leaf Miners  
Leaf Rollers  
Leaf Tiers  
Leafhoppers  
Lesser Appleworms  
Lesser Peach Tree Borers  
Loopers  
Lygus Bugs  
Mexican Bean Beetles  
Mites  
Mole Crickets  
Naval Orangeworms  
Northern Pine Weevils  
Oriental Fruit Moths  
Painted Lady Caterpillars  
Pea Weevils  
Peach Tree Borers  
Peach Twig Borers

Pear Psyllid  
Pear Slugs  
Pecan Leaf Phylloxera  
Pecan Nut Casebearers  
Pecan Spittlebugs  
Pecan Stem Phylloxera  
Pecan Weevils  
Pepper Weevils  
Periodical Cicadas  
Pickleworms  
Pillbugs  
Pine Chafers  
Pine Coreid Bugs  
Plant Bugs  
Plum Curculios  
Red Pine Sawflies  
Red-striped Fireworms  
Redheaded Pine Sawflies  
Rindworms  
Salt Marsh Caterpillars  
Sap Beetles  
Scales  
Sod Webworms  
Sowbugs  
Spiders  
Spittlebugs  
Stink Bugs  
Tarnished Plant Bugs  
Tent Caterpillars  
Thrips  
Tobacco Hornworms  
Tufted Apple Budmoths  
Velvet Bean Caterpillars  
Walnut Husk Flies  
Western Bean Cutworms  
Whiteflies

### FLYING INSECTS INCLUDING:

Blow Flies  
Chiggers  
Clothes Moths  
Crane Flies  
Deer Flies  
Face Flies

## General information:

Do not apply this product in patient rooms or in any rooms while occupied by the elderly or infirm. Do not apply to classrooms while in use. Do not apply in institutions (including libraries, schools, sports facilities, etc.) in the immediate area where occupants are present.

Onslaught® Microencapsulated Insecticide is a microencapsulated suspension concentrate containing 0.54 pounds active ingredient per gallon.

Onslaught® Microencapsulated Insecticide is designed to provide long residual activity against pests listed on this label when applied as surface or spot treatments, injected into wall voids, or as a crack and crevice spray in and around residential and commercial structures, as well as on turf and landscaping. Onslaught® Microencapsulated Insecticide may be sprayed on any surface that will not be stained by water. DO NOT use this product as a fogger or apply it as a space spray. Do not apply in electrical conduits, junction and switch boxes, motor housings, or other electrical equipment due to shock hazard from water-based spray.

Use only in areas described. Remove pets and cover fish aquariums and pets' food and water dishes before spraying. Keep all people (especially children) and pets out of areas being treated and restrict access to these areas until all surfaces are dry. DO NOT leave product where children or pets can come into contact with it. DO NOT allow spray to contact food or food-containing surfaces, feed, or water supplies. Thoroughly wash dishes and food handling utensils with soap and water if they become contaminated with this product. Food utensils such as teaspoons and measuring cups must not be used for food purposes after use with pesticides.

## Application in food processing/handling establishments:

Application is permitted within food and non-food areas of food service and handling establishments (places other than private residences) including but not limited to: restaurants, meat processing plants, grocery stores, bakeries, food manufacturing and processing establishments, and food warehousing establishments. Do not treat establishments where livestock feed is present.

Food areas include areas for receiving, storing, packing, canning, bottling, wrapping, boxing, preparing, edible waste storage, and enclosed processing systems, mills, and dairies. Serving areas are places where prepared foods are served, such as dining rooms, but excluding areas where foods may be prepared or held. Non-food areas include garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, garages, mop closets, and storage after canning or bottling.

General surface applications: Do not apply Onslaught® Microencapsulated Insecticide as a general surface application in food areas of the facility when the facility is in operation or food is exposed. Do not apply directly to food. Cover or remove all food and food processing equipment prior to application. After spraying in meat packing plants, bakeries, and other food processing plants, wash with an effective cleaning compound, and then rinse water all equipment, benches, shelving, etc. where exposed food will be handled with potable water. Repeat applications as needed, but do not exceed more than one (1) application every (14) days.

Spot or crack and crevice applications: Spot or crack and crevice applications may be made while facility is in operation, provided exposed food is covered or removed from the area being treated. Do not apply directly to food or food-handling surfaces. Repeat applications as needed, but do not exceed more than one (1) application every (14) days.

In the home, cover exposed food and do not allow spray to contact food/feed surfaces. If spray does contact these surfaces, clean surfaces with soap and water.

## MIXING INSTRUCTIONS

Onslaught® Microencapsulated Insecticide should be mixed with water and applied with hand pressurized or power operated sprayers. Shake container of Onslaught® Microencapsulated Insecticide before diluting. Clean spray equipment before use. For dilution, add approximately half the required water to spray tank and then add the appropriate amount of Onslaught® Microencapsulated Insecticide. Agitate and slowly add the remaining water. Agitate spray thoroughly before using and also occasionally during use to ensure dispersion. If spray filter screens are used, they should be 50 mesh or larger. Use 0.5 fluid ounces (15 cc.) of Onslaught® Microencapsulated Insecticide in 1 gallon of water to make a spray mix containing 0.025% active ingredient. Use 1.0 fluid ounce (30 cc.) per gallon of water to make a spray containing 0.05% active ingredient. Use 0.025% solution for light infestations or as a maintenance control rate. Use 0.05% for heavy infestations or as an initial clean out rate. Apply two-second bursts of spray per square foot of area being treated. Avoid excessive application. Dampen surfaces but not to the point of saturation or run-off. Only dilute Onslaught® Microencapsulated Insecticide with water.

## Formula for determining the active ingredient content of the finished spray mixture:

The following formula may be used to determine the percent active ingredient that is in the sprayer tank after mixing Onslaught® Microencapsulated Insecticide:

$$\frac{(6.4) \times (\text{number of fl. oz. of concentrate added to tank})}{(\text{Gallons of finished spray mix}) \times (128)} = \% \text{ active ingredient of spray mix}$$

## Tank mixing:

Onslaught® Microencapsulated Insecticide may be tank mixed with an insect growth regulator such as NyGuard® IGR Concentrate or pyrethrum-containing products or any other currently registered pesticides unless expressly prohibited by the product label. The resulting tank mix may be applied in areas where these products are allowed to be sprayed. Do not tank mix Onslaught® Microencapsulated Insecticide with products containing dichlorvos (DDVP).

## INDOOR USE

### Application rates for indoor structural pests:

Pests	Concentration of active ingredient	Dilution rate in water	Application rate of mixed solution	Application method(s)
<b>Crawling insects</b> Light infestation Heavy infestation	0.025% 0.050%	0.5 fl. oz./gallon 1.0 fl. oz./gallon	1 gallon/1,000 sq. ft.	Apply as a coarse, wet spray • Broadcast surface spray • Crack and crevice treatment
<b>Flying insects</b>	0.025%	0.5 fl. oz./gallon	1 gallon/1,000 sq. ft.	• Spot treatment • Injected into wall voids
<b>Stinging insects</b>	0.050%	1.0 fl. oz./gallon	1 gallon/1,000 sq. ft.	
<b>Ticks</b>	0.025%	0.5 fl. oz./gallon	1 gallon/1,000 sq. ft.	
<b>Fleas, Lice and Bed Bugs</b>	0.025%	0.5 fl. oz./gallon	1 gallon/1,000 sq. ft.	Apply as noted above, with a fine particle spray

### Crawling insect pests indoors:

Apply as a coarse, wet spray to surfaces where these pests are normally found. Treat floors, baseboards, around doors and windows, in attics, crawl spaces, eaves, corners, closets, walls, utility pipes, storage areas, and all cracks and crevices. Treat underneath sinks, dishwashers, refrigerators, stoves, the underside of shelves, drawers, cabinets, areas behind pipes, and in all places where these insects shelter. Contact as many insects as possible with direct spray. Repeat applications as needed, but do not exceed more than one (1) application every seven (7) days.

### For Ant control indoors:

Apply to ant trails, around garbage receptacles, and near food sources. Also apply around doors and windows and wherever these pests may find entrance to the structure. Remove sources of food through basic hygiene practices whenever possible.

### Flying insect pests indoors:

Treat insect resting areas such as walls and ceilings, screens, around windows, doors, and light fixtures, and other surfaces that attract flying insects. Initiate treatments at the beginning of fly season, and repeat treatments during periods of heavy infestation.

### Stinging insect pests indoors:

Apply spray to nests late in the evening when stinging insects are at rest. Thoroughly spray nest, nest entrance, and surrounding areas where insects land or walk. Treat around doors and windows, in attics, crawl spaces, and possible harborage sites or points of entry.

### For control of Brown Dog Ticks:

Thoroughly apply as a spot treatment to infested areas such as pet beds and resting areas, nearby cracks and crevices, along and behind baseboards, window and door frames, and localized areas of floor and floor coverings where these pests may be present. DO NOT SPRAY PETS WITH THIS PRODUCT. Treat dogs and cats with a product registered for use on animals.

### For control of Fleas:

Thoroughly apply as a fine particle broadcast spray to infested rugs, carpets, and pet resting areas. Prior to treatment, aquariums and fish bowls should be covered, and pet animals should be removed from the area being treated. Do not permit humans or pets to contact treated surfaces until spray has dried. Old pet bedding should be removed and replaced with clean, fresh bedding after treatment. DO NOT SPRAY PETS WITH THIS PRODUCT. Treat dogs and cats with a product registered for use on animals.

### To control Bed Bugs:

Thoroughly clean and sanitize mattresses and box springs. Treat mattresses and box springs with an approved pesticide, such as a pyrethrin aerosol. Apply Onslaught® Microencapsulated Insecticide as a spot treatment to potential harborage sites and migration paths, and cracks and crevices, around baseboards, floorboards, head-boards, and walls.

### For control of stored product pests:

Spray thoroughly around and into floor drains, non-food conveyors, benches, pipes, pallets, moist areas, storage racks, pieces of equipment, and other areas where stored product pests may be found. Tank-mix or sequential use of an insect growth regulator, such as NyGuard® IGR Concentrate, is recommended to break the insect reproduction cycle. Do not apply this spray to surfaces or utensils that may come in contact with food, since excessive residues in food may result.

**To treat voids in equipment and structures:**

To kill insects harboring in wall voids and other inaccessible spaces in equipment and structures, use injection equipment designed for deep void applications. Follow the injection equipment manufacturer's recommendations for proper set up and air pressures. Place the applicator tip at or into the void space to be treated. For inaccessible voids, it may be necessary to drill an access hole(s). Inject product into the void space in short bursts, allowing air pressure to push insecticide deep into the space. Avoid applying to the point of runoff or drip. Ventilate area thoroughly before re-entry. Do not reapply more than every fourteen (14) days.

**To kill the accessible stages of listed granary insects:**

Pest management professionals and grain producers may use this concentrate to treat grain storage facilities, and other listed areas, for stored product pest control. For control of exposed adult and immature stages of stored product pests, apply to cracks, crevices, and other surfaces where the pests have been seen or have harborage. Treat areas where products are stored before filling with the product. Apply at the rates listed above, using one gallon of spray mix per 1,000 square feet of surface area to be treated. Cleaning all areas prior to use will increase levels of control. Any foodstuffs infested with pests should be removed and destroyed. Do not apply when food-processing facility is in operation or foods are exposed. Do not apply this spray to surfaces or utensils that may come into contact with food.

Repeat application as needed, but DO NOT exceed more than one application every fourteen (14) days.

**USE IN AND AROUND LIVESTOCK HOUSING**

For use in and around unoccupied areas of livestock facilities, such as, but not limited to: barns, cow and calf pens and hutches, dairy barns and milk rooms, hog barns, horse barns, sheep barns, poultry houses, and rabbit hutches. Do not contaminate milk, food, or drinking water. Remove animals from area being treated. Cover feeders and waterers. Do not apply or allow insecticide to drift onto animals. Do not allow animals to enter treated areas until spray solution has dried.

Pests	Concentration of active ingredient	Dilution rate in water	Application method(s)
<b>Crawling insects</b> * Including Litter Beetles	0.050%	1.0 fl. oz./gallon or 1.0 fl. oz. in sufficient water to cover 1,000 sq. ft.	Apply as a coarse, wet spray • General surface spray • Crack and crevice treatment
<b>Flying insects</b>	0.025%	0.5 fl. oz./gallon or 0.5 fl. oz. in sufficient water to cover 1,000 sq. ft.	• Spot treatment • Injected into wall voids

**Crawling insect pests in livestock and poultry premises:**

Apply as a general surface, spot treatment, and/or crack and crevice treatment. Apply to floors and vertical and overhead surfaces where crawling insects are or may be present. Treat stanchions, pipes, windows, doors, posts, cage framing, gates, under (but not in) feeders, and other areas where insects hide or congregate. To reduce immigration of insects, make a perimeter treatment around the outside of building foundations. Apply in a uniform band 1 to 3 feet up the exterior foundation wall and 3 to 6 feet out from the foundation.

**Litter Beetle control in livestock and poultry premises:**

To control Litter Beetles (darkling, hide, and carrion beetles), apply Onslaught® Microencapsulated Insecticide to walls and floors at cleanout and before reintroduction of animals. Treat areas where beetles frequently occur, such as walls, supports, cages, cage framing, stalls, and around feeders. To reduce immigration of insects, make a perimeter treatment around the outside of building foundations. Apply in a uniform band 1 to 3 feet up the exterior foundation wall and 3 to 6 feet out from the foundation.

**Flying insect pests in livestock and poultry premises:**

For residual control of flying insects, treat insect resting areas such walls, ceilings, screens, around windows, doors, light fixtures, and other surfaces that attract flying insects. Initiate treatments at the beginning of fly season and repeat treatments during periods of heavy infestation.

**OUTDOOR USE**

Do not spray in or near fish ponds or other bodies of water.

Not for use on plants being grown for sale or other commercial use. Not for use in nurseries, sod farms or golf courses.

**Application rates for outdoor pests:**

	Concentration of active ingredient	Dilution rate in water	Application method(s)
<b>Treating exterior walls, foundations and structures</b>	0.025% to 0.050%	0.5 to 1.0 fl. oz./gallon or 0.5 to 1.0 fl. oz. in sufficient water to cover 1,000 sq. ft.	Apply as a coarse, wet spray • General surface spray • Crack and crevice treatment • Spot treatment
<b>Treating nests and harborage areas of stinging insects</b>	0.050%	1.0 fl. oz./gallon or 1.0 fl. oz. in sufficient water to cover 1,000 sq. ft.	
<b>To control Swarming Termites and wood destroying pests</b>	0.050%	1.0 fl. oz./gallon or 1.0 fl. oz. in sufficient water to cover 1,000 sq. ft.	
<b>Treating ornamental trees and landscapes</b>	0.005% to 0.050%	0.1 to 1.0 fl. oz./gallon or 0.1 to 1.0 fl. oz. in sufficient water to cover 1,000 sq. ft.	Apply as a coarse, wet spray • Broadcast spray treatment • Foliar spray treatment
<b>Mosquito breeding sites</b>	0.025%	0.5 fl. oz./gallon or 0.5 fl. oz. in sufficient water to cover 1,000 sq. ft.	
<b>Treating lawns and turf</b>	0.005% to 0.050%	0.1 to 1.0 fl. oz./gallon or 0.1 to 1.0 fl. oz. in sufficient water to cover 1,000 sq. ft. or 4.3 to 43 fl. oz. per acre	Apply as a coarse, wet spray • Broadcast spray treatment

**Treating exterior walls, foundations and structures:**

To control infestations of listed pests, treat exterior surfaces of buildings, walls, window frames, around garbage cans, eaves, cracks and crevices, porches, decks, gazebos, patios, carports, garages, fence lines, and other areas where pests are active or may be hiding.

To help prevent insect infestation of buildings: Treat a 2 to 6 foot band of soil or other substrate adjacent to buildings. Treat building foundation to a height of 2 to 3 feet. Treat exterior walls, eaves, cracks and crevices, and other areas where listed pests are active, and may find entrance into building.

Apply with sufficient water to adequately cover the area being treated, but do not allow dripping or run-off to occur. Alternate mixing directions are to use 2 to 4 fluid ounces of concentrate per 50 gallons of water.

**Stinging insect control outdoors:**

For stinging insects, apply spray solution to nests and harborage areas late in the evening when stinging insects are commonly at rest. Thoroughly spray nest, nest entrance, harborage sites, and surrounding areas where insects land or walk. For nests inside walls, inject sufficient spray to treat nest. Do not use in areas where an electrical shock hazard exists. For applications made when pests are active, applicator should wear protective equipment as required. Repeat application when there are signs of renewed insect activity.

For Yellow Jacket control, Onslaught® Microencapsulated Insecticide can be mixed with baits in traps. Follow trap instructions for preparation of bait.

For control of Fire Ants, combine broadcast application with mound drenches to control foraging workers and newly mated fly-in queens. Apply Onslaught® Microencapsulated Insecticide as a broadcast application at a rate of 8 fl. oz. per 1,000 square feet. Treat mounds with a registered fire ant mound treatment.

**Insect control in ornamental trees and landscapes:**

For residential and commercial trees, shrubs, ground covers, and bedding and foliage plants that will not be harvested for food, apply Onslaught® Microencapsulated Insecticide in appropriate volumes of water to obtain thorough coverage. Apply as a full-coverage foliar spray, applying to the point of drip but not runoff. Treat active soil pests with an application to surrounding soil. Repeat treatments as necessary to achieve control, using higher application rates as pest pressure and foliage area increases. Repeat applications as necessary, but no more than once every seven (7) days. Certain plants may be sensitive to the final spray solution. A small-scale test is recommended to verify safety to ornamental plants. Spray and observe for one (1) week prior to application of an entire planting.

**Insect control on lawns, turf grass and turf:**

For best results, lawn or turf grass should be mowed 1 to 2 days before spraying. Treat with spraying equipment or a hand sprayer. Use application volumes of up to 10 gallons per 1,000 square feet to get uniform coverage when treating dense grass foliage. For low volume applications using less than 2 gallons of spray solution per 1,000 square feet, immediate irrigation with at least 0.25 inches of water is recommended to improve effectiveness on sub-surface pests.

When hand spraying, spray using a slow, even sweeping motion, making sure to cover the entire lawn or turf grass area where pests are observed. Spray under ornamentals and trees. Repeat treatments may be necessary at 7 to 14 day intervals.

**Rate range:**

Use lower rate range for pests that are commonly exposed and will be contacted by spray solution at the time of application. For pests that will not be contacted by spray solution at the time of application, use the upper rate range. Use 1 fl. oz. per 1,000 sq. ft. when treating Mole Crickets and Chinch Bugs.

**To kill Swarming Termites: OUTDOORS ONLY.**

Apply spray mix as a coarse, wetting spray when Swarming Termites are seen emerging from woodpiles, wooden fence posts, wooden structures, or from the ground. Swarming usually occurs in the spring or at other times when a termite colony becomes overcrowded and new reproductive termites with wings emerge and fly away to mate and establish new colonies. This treatment will control the sprayed termites and will not protect the structure from which the swarm is coming. Use only as a contact spray to kill emerging reproductive (winged) and worker termites emerging from infested wood. This treatment is not a substitute for a comprehensive termite control program.

**To kill wood destroying pests: OUTDOORS ONLY.**

Apply spray mix as a coarse, wetting spray to exposed pests and to the damaged areas of wood, spraying into galleries or tunnels in the exposed wood. Also spray around doors, window and door frames, and other areas where these pests may hide or enter the house. Spray into cracks and crevices and, if necessary, drill small holes and spray into inaccessible wooden structural voids where these pests are suspected.

**Mosquito breeding sites:**

Mosquito populations may be reduced by application of Onslaught® Microencapsulated Insecticide to sites where Mosquitoes rest, harbor, and breed. Apply spray solution into tall grass, shrubbery, and around backyards and lawns where these pests may hover or rest. Apply while air is still.

Not for wide area mosquito use. Do not apply with hand held or truck mounted cold aerosol ULV sprayers and thermal fogging devices.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store in a cool, dry, well-ventilated area. Keep container closed.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. In case of spill or leak, soak up with sand, earth, or synthetic absorbents. Do not use alkaline absorbents or clean spill area with alkaline detergents.

**CONTAINER DISPOSAL:** Do not reuse empty container. Wrap container in several layers of newspaper and discard in trash.



Onslaught, NyGuard, McLaughlin Gormley King Company and MGK are registered trademarks of McLaughlin Gormley King Company. Manufactured by McLaughlin Gormley King Company®. ©2009 McLaughlin Gormley King Company. All rights reserved. 8810 Tenth Avenue North, Minneapolis, MN 55427 037-1678a/5M-0609

PRECAUTIONARY STATEMENTS  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION**

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wear protective eyewear (goggles, face shield or safety glasses with side shields). Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish and other aquatic invertebrates. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirement of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product into sewer systems without previously notifying sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

**PHYSICAL OR CHEMICAL HAZARDS**

Do not use or store near heat or open flame. Do not use this product in or on electrical equipment due to the possibility of shock hazard.

**FIRST AID**

**IF SWALLOWED:**

- Call a poison control center or doctor immediately for treatment advice.
- Do not give **any** liquid to the person.
- Do not induce vomiting unless told to do so by a poison control center or a doctor.
- Do not give anything by mouth to an unconscious person.

**IF IN EYES:**

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first five minutes, then continue rinsing eyes.
- Call a poison control center or doctor for treatment advice.

**IF ON SKIN**

**OR CLOTHING:**

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

**NOTE TO PHYSICIAN:**

Contains petroleum distillate—vomiting may cause aspiration pneumonia.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For information regarding medical emergencies or pesticide incidents, call 1-888-740-8712.